

Recombinant Human PLCB Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP50796

Size: 10ug, 100ug

Protein Name: 1-acyl-sn-glycerol-3-phosphate acyltransferase beta (EC 2.3.1.51) (1-acylglycerol-3-phosphate O-acyltransferase 2) (1-AGP acyltransferase 2) (1-AGPAT 2) (Lysophosphatidic acid acyltransferase beta) (LPAAT-beta)

Synonym: AGPAT2

UniProt ID: O15120

UniProt Link: www.uniprot.org/uniprotkb/O15120/entry

Species Source: Human

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 278 AA

Molecular Weight: 31 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MELWPCLA AALLLLLLLVQLSRAAEFYAKVALYCALCFTVSAVASLVCLLRHGGRTVENMSIIGWFVR
SFKYFYGLRFEVRDPRRLQEARPCVIVSNHQSILDMGLMEVLPERCVQIAKRELLFLGPVGLIMYLG
GVFFINRQRSSTAMTVMADLGERMVRENKLVWIYPEGTRNDNGDLLPFKKGAFYLAVQAQVPIVPV
VYSSFSSFYNTKKKFFTSGTVTVQVLEAIPTSGLTAADV PALVDTCHRAMRTTFLHISKTPQENGATA
GSGVQPAQ

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant 1-acyl-sn-glycerol-3-phosphate acyltransferase beta (EC 2.3.1.51) (1-acylglycerol-3-phosphate O-acyltransferase 2) (1-AGP acyltransferase 2) (1-AGPAT 2) (Lysophosphatidic acid acyltransferase beta) (LPAAT-beta) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Converts 1-acyl-sn-glycerol-3-phosphate (lysophosphatidic acid or LPA) into 1,2-diacyl-sn-glycerol-3-phosphate (phosphatidic acid or PA) by incorporating an acyl moiety at the sn-2 position of the glycerol backbone.

Subcellular Localization: Endoplasmic reticulum membrane ; Multi-pass membrane protein .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.