

Recombinant Mouse TAF9 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51813

Size: 10ug, 100ug

Protein Name: Transcription initiation factor TFIID subunit 9 (RNA polymerase II TBP-associated factor subunit G) (Transcription initiation factor TFIID 31 kDa subunit) (TAFII-31) (TAFII31) (Transcription initiation factor TFIID 32 kDa subunit) (TAFII-32) (TAFII32)

Synonym: Taf9, Taf2g

UniProt ID: Q8VI33

UniProt Link: www.uniprot.org/uniprotkb/Q8VI33/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 264 AA

Molecular Weight: 29 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MESGKMASPKSMPKDAQMMAQILKDMGITEYEPRVINQMLEFAFRYVTTILDDAKIYSSHAKKATVD
ADDVRLAIQCRADQSFTSPPPRDFLLDIARQRNQTPPLIKPYSGPRLPPDRYCLTAPNYRLKSLQKK
APAPAGRITVPRLSVGSVSSRPSTPTLGTPTQTMSVSTKVGTPMSLTGQRFTVQMPASQSPAVKA
SIPATSTVQNVLINPSLIGSKNILITTNMVSQNTAESANALKRKREDDDDDDDDDDDDDDDDYDNM

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Transcription initiation factor TFIID subunit 9 (RNA polymerase II TBP-associated factor subunit G) (Transcription initiation factor TFIID 31 kDa subunit) (TAFII-31) (TAFII31) (Transcription initiation factor TFIID 32 kDa subunit) (TAFII-32) (TAFII32) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: The TFIID basal transcription factor complex plays a major role in the initiation of RNA polymerase II (Pol II)-dependent transcription. TFIID recognizes and binds promoters with or without a TATA box via its subunit TBP, a TATA-box-binding protein, and promotes assembly of the pre-initiation complex (PIC). The TFIID complex consists of TBP and TBP-associated factors (TAFs), including TAF1, TAF2, TAF3, TAF4, TAF5, TAF6, TAF7, TAF8, TAF9, TAF10, TAF11, TAF12 and TAF13. TAF9 is also a component of the TBP-free TAFII complex (TFTC), the PCAF histone acetylase complex and the STAGA transcription coactivator-HAT complex. TAF9 and its paralog TAF9B are involved in transcriptional activation as well as repression of distinct but overlapping sets of genes. Essential for cell viability. May have a role in gene regulation associated with apoptosis.

Subcellular Localization: Nucleus .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.