

Recombinant Mouse IFIT2 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51673

Size: 10ug, 100ug

Protein Name: Interferon-induced protein with tetratricopeptide repeats 2 (IFIT-2) (Glucocorticoid-attenuated response gene 39 protein) (GARG-39) (Interferon-induced 54 kDa protein) (IFI-54K) (P54)

Synonym: Ifit2, Garg39, Ifi54, Isg54

UniProt ID: Q64112

UniProt Link: www.uniprot.org/uniprotkb/Q64112/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 472 AA

Molecular Weight: 55 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MSTTSKESLVCNLRQLKCHFTWNLIAEDESDEFEDRVFNKDEFQNSEFKATMCNILAYVKHCRGLN
EAALQCLGEAEGFIQQHPDQVEIRSLVTWGNyawvyyHMGQFSKAQAYLDKVKQVCKKFSSPYRI
ENPALDCEEGWARLKCTKNQNERVKVCFQKALEKDPKNPEFTSGWAIAFYRLDDWPARNYCIDSLE
QAIQLSPDNTYVKVLLALKLDAVHVHKNQAMALVEEALKKDPSAIDTLLRAARFYCKVYDTRAIQLL
RKALEKLPPNAYVHYHMGCCYRSKVHMLNRREMFSGDRKKLEELIQLAVNHLRKAEEIKEMLEY
SCSFLADLYIIAKKYDEADYYFQKELSKDLPPGPKQLLHLRYGNFQFFQMKRQDKAIHYHMEGVKIKK
KTIPQKKMREKLQRIALRRLHEDESSEALHILAFLQENGGGQQADKDSERGVDSANQVPSASLDED
GAEY

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Interferon-induced protein with tetratricopeptide repeats 2 (IFIT-2) (Glucocorticoid-attenuated response gene 39 protein) (GARG-39) (Interferon-induced 54 kDa protein) (IFI-54K) (P54) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: IFN-induced antiviral protein which inhibits expression of viral messenger RNAs lacking 2'-O-methylation of the 5' cap. The ribose 2'-O-methylation would provide a molecular signature to distinguish between self and non-self mRNAs by the host during viral infection. Viruses evolved several ways to evade this restriction system such as encoding their own 2'-O-methylase for their mRNAs or by stealing host cap containing the 2'-O-methylation (cap snatching mechanism). Binds AU-rich viral RNAs, with or without 5' triphosphorylation, RNA-binding is required for antiviral activity. Can promote apoptosis.

Subcellular Localization: Cytoplasm . Endoplasmic reticulum .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.