

Recombinant Mouse PD1L1 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51886

Size: 10ug, 100ug

Protein Name: Programmed cell death 1 ligand 1 (PD-L1) (PDCD1 ligand 1) (Programmed death ligand 1) (B7 homolog 1) (B7-H1) (CD antigen CD274)

Synonym: Cd274, B7h1, Pdcd1l1, Pdcd1lg1, Pdl1

UniProt ID: Q9EP73

UniProt Link: www.uniprot.org/uniprotkb/Q9EP73/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 290 AA

Molecular Weight: 33 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MRIFAGIIIFTACCHLLRAFTITAPKDLYVVEYGSNVTMECRFPVERELDLLALVVYWEKEDEQVIQFVA
GEEDLKPKQHSNFRGRASLPKDQLLKGNAAALQITDVKLQDAGVYCCIIISYGGADYKRITLKVNPYRKI
NQRISVDPATSEHELICQAEGYPEAEVIWTNSDHQPVSGKRSVTTSRTEGMLLNVTSSLRVNATAND
VFYCTFWRSQPGQNHTAELIPELPPATHPPQNRTHWVLLGSILLFLIVVSTVLLFLRKQVRMLDVEKC
GVEDTSSKNRNDTQFEET

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant Programmed cell death 1 ligand 1 (PD-L1) (PDCD1 ligand 1) (Programmed death ligand 1) (B7 homolog 1) (B7-H1) (CD antigen CD274) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Plays a critical role in induction and maintenance of immune tolerance to self (PubMed:11238124). As a ligand for the inhibitory receptor PDCD1/PD-1, modulates the activation threshold of T-cells and limits T-cell effector response (PubMed:11238124). Through a yet unknown activating receptor, may costimulate T-cell subsets that predominantly produce interleukin-10 (IL10) (PubMed:11015443, PubMed:12719480).; The PDCD1-mediated inhibitory pathway is exploited by tumors to attenuate anti-tumor immunity and escape destruction by the immune system, thereby facilitating tumor survival (PubMed:12218188, PubMed:27281199). The interaction with PDCD1/PD-1 inhibits cytotoxic T lymphocytes (CTLs) effector function (PubMed:12218188). The blockage of the PDCD1-mediated pathway results in the reversal of the exhausted T-cell phenotype and the normalization of the anti-tumor response, providing a rationale for cancer immunotherapy (PubMed:12218188).

Subcellular Localization: Cell membrane ; Single-pass type I membrane protein . Early endosome membrane ; Single-pass type I membrane protein . Recycling endosome membrane ; Single-pass type I membrane protein .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.