

Recombinant Mouse RSAD2 Protein

For Research Use Only

1. Basic Product Information

Catalog No.: REP51759

Size: 10ug, 100ug

Protein Name: S-adenosylmethionine-dependent nucleotide dehydratase RSAD2 (SAND) (EC 4.2.-.-) (Radical S-adenosyl methionine domain-containing protein 2) (Virus inhibitory protein, endoplasmic reticulum-associated, interferon-inducible) (Viperin)

Synonym: Rsad2, Vig1

UniProt ID: Q8CBB9

UniProt Link: www.uniprot.org/uniprotkb/Q8CBB9/entry

Species Source: Mouse

Expression System: HEK293T eukaryotic cell expression system Sequence

Length: 362 AA

Molecular Weight: 42 KD

Protein Tag: C-terminal His6

2. Amino Acid Sequence

MGMLVPTALAAARLLSLFQQQLGSLWSGLAILFCWLRIALGWLDPGKEQPQVRGEPEDTQETQEDGN
STQPTTPVSVNYHFTRQCNYKCGFCFHTAKTSFVLPLEEAKRGLLLLKQAGLEKINFSGGEPFLQDR
GEYLGKLVRFCKEELALPSVSIVSNGSLIRERWFKDYGEYLDILAISCDSEVQVNALIGRGQGKKNH
VENLQKLRRWCRDYKVAFKINSVINRFNVEDMNEHIKALSPVRWKVFQCLLIEGENSGEDALREAE
RFLISNEEFETFLERHKEVSCLVPESNQKMKDSYLILDEYMRFLNCTGGRKDPSKSILDVGVEEAIKF
SGFDEKMFLKRGGKYVWSKADLKLDW

3. Buffer Composition

Liquid format: PBS-based buffer containing 50% glycerol. Glycerol concentration can be customized upon customer request; please state special requirements when placing your order.

Lyophilized format: Buffer before lyophilization is PBS supplemented with 6% trehalose.

4. Storage Conditions

Upon receipt, store the product at -20 °C or -80 °C. Aliquot the product for on-demand use to preserve protein activity and prevent denaturation and inactivation caused by repeated freeze-thaw cycles.

5. Product Description

This product is recombinant S-adenosylmethionine-dependent nucleotide dehydratase RSAD2 (SAND) (EC 4.2.-.-) (Radical S-adenosyl methionine domain-containing protein 2) (Virus inhibitory protein, endoplasmic reticulum-associated, interferon-inducible) (Viperin) produced using the HEK293T eukaryotic cell expression system. It is suitable for in-vitro functional assays, protein-protein interaction studies, antibody preparation and other biomedical research applications.

Protein Function: Interferon-inducible antiviral protein which plays a major role in the cell antiviral state induced by type I and type II interferon. Catalyzes the conversion of cytidine triphosphate (CTP) to 3'-deoxy-3',4'-didehydro-CTP (ddhCTP) via a SAM-dependent radical mechanism. In turn, ddhCTP acts as a chain terminator for the RNA-dependent RNA polymerases from multiple viruses and directly inhibits viral replication. Therefore, inhibits a wide range of DNA and RNA viruses (By similarity). Also promotes TLR7 and TLR9-dependent production of IFN-beta production in plasmacytoid dendritic cells (pDCs) by facilitating 'Lys-63'-linked ubiquitination of IRAK1 by TRAF6. Plays a role in CD4+ T-cells activation and differentiation. Facilitates T-cell receptor (TCR)-mediated GATA3 activation and optimal T-helper 2 (Th2) cytokine production by modulating NFkB1 and JUNB activities. Can inhibit secretion of soluble proteins (By similarity) (PubMed:17686841, PubMed:19047684, PubMed:21435586, PubMed:21880757).

Subcellular Localization: Endoplasmic reticulum membrane ; Peripheral membrane protein ; Cytoplasmic side . Golgi apparatus . Endoplasmic reticulum . Lipid droplet . Mitochondrion . Mitochondrion inner membrane . Mitochondrion outer membrane .

6. Notes

- Avoid repeated freeze-thaw cycles to maintain protein stability and biological activity.
- Please specify special buffer-composition requirements at the time of order placement.
- For research use only. Not intended for clinical diagnosis, therapeutic use or commercial manufacturing.